



Taking off personal protective equipment (PPE) for non-aerosol generating procedures (AGPs)*


Please see donning and doffing video to support this guidance: https://youtu.be/-GncQ_ed-9w

• PPE should be removed in an order that minimises the risk of self-contamination

• Gloves, aprons (and eye protection if used) should be taken off in the patient's room or cohort area

1 Remove gloves. Grasp the outside of glove with the opposite gloved hand; peel off.

Hold the removed glove in the remaining gloved hand.



Slide the fingers of the un-gloved hand under the remaining glove at the wrist.

Peel the remaining glove off over the first glove and discard.



2 Clean hands.



3 Apron.
Unfasten or break apron ties at the neck and let the apron fold down on itself.



Break ties at waist and fold apron in on itself – do not touch the outside – **this will be contaminated.** Discard.



4 Remove eye protection if worn.

Use both hands to handle the straps by pulling away from face and discard.



5 Clean hands.



6 Remove facemask once your clinical work is completed.



Untie or break bottom ties, followed by top ties or elastic, and remove by handling the ties only. Lean forward slightly. Discard. **DO NOT** reuse once removed.

7 Clean hands with soap and water.



*For the PPE guide for AGPS please see:

www.gov.uk/government/publications/covid-19-personal-protective-equipment-use-for-aerosol-generating-procedures